




1. nonionic surfactants derived from saturated and/or unsaturated primary, secondary, and/or branched, amine, amide, amine-oxide, fatty alcohol, fatty acid, alkyl phenol, and/or alkyl aryl carboxylic acid compounds having from about 6 to about 22 carbon atoms in a hydrophobic chain, wherein at least one active hydrogen of said compounds is ethoxylated with ≤ 50 ethylene oxide moieties to provide an HLB of from about 8 to about 20;
2. nonionic surfactants with bulky head groups selected from:
 - a. surfactants having the formulas:




wherein Y'' = N or O; and each R⁵ is selected independently from the following: -H, -OH, -(CH₂)_xCH₃, -O(OR²)_z-H, -OR¹, -OC(O)R¹, and -CH(CH₂-(OR²)_{z'}-H)-CH₂-(OR²)_z-C(O)R¹, wherein R¹ is selected from the group consisting of saturated or unsaturated, primary, secondary or branched chain alkyl or alkyl-aryl hydrocarbons; said hydrocarbon chain having a length of from about 6 to about 22, wherein each R² is selected from the following groups or combinations of the following groups: -(CH₂)_n- and/or -[CH(CH₃)CH₂]- wherein n is from 1 to 4; and wherein x is from 0 to about 3, and z, z', and z'' are from about 5 to about 20;

- b. polyhydroxy fatty acid amide surfactants of the formula:



wherein: each R¹ is H, C₁-C₄ hydrocarbyl, C₁-C₄ alkoxyalkyl, or hydroxyalkyl; R² is a C₅-C₂₁ hydrocarbyl moiety; and each Z is a polyhydroxyhydrocarbyl moiety having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an ethoxylated derivative thereof;


- c. surfactants having the formula



wherein R¹ is selected from the group consisting of saturated or unsaturated, primary, secondary or branched chain alkyl or alkyl-aryl hydrocarbons; said hydrocarbon chain having a length of from about 6 to about 22; Y' is selected from the following groups: -O-; -N(A)-; and mixtures thereof; and A is selected from the following groups: H; R¹; -(R²-O)_z-H; -(CH₂)_xCH₃; phenyl, or substituted aryl,

wherein x is from 0 to about 3 and total z is from about 5 to about 30; each R^2 is selected from the following groups or combinations of the following groups: $-(CH_2)_n-$ wherein n is from about 1 to about 4 and/or $-[CH(CH_3)CH_2]-$; each R^5 is selected from the following groups: $-OH$; and $-O(R^2O)_x-H$; and m is from about 2 to about 4; and

- d. mixtures thereof;
3. surfactant complexes formed by one surfactant ion being neutralized with surfactant ion of opposite charge or an electrolyte ion that is suitable for reducing dilution viscosity;
4. block copolymer surfactants comprising polyethylene oxide moieties and propylene oxide moieties;
5. cationic surfactants having the formula:



wherein R^1 is selected from the group consisting of saturated or unsaturated, primary, secondary or branched chain alkyl or alkyl-aryl hydrocarbons; said hydrocarbon chain having from about 6 to about 22 carbon atoms; each R^2 is selected from the following groups or combinations of the following groups: $-(CH_2)_n-$ and/or $-[CH(CH_3)CH_2]-$; Y is selected from the following groups: $=N^+-(A)_q$; $-(CH_2)_n-N^+-(A)_q$; $-B-(CH_2)_n-N^+-(A)_2$; $-(phenyl)-N^+-(A)_q$; $-(B-phenyl)-N^+-(A)_q$; with n being from about 1 to about 4, wherein each A is independently selected from the following groups: H; C_{1-5} alkyl; R^1 ; $-(R^2O)_x-H$; $-(CH_2)_xCH_3$; phenyl, and substituted aryl; where x is from 0 to about 3; and each B is selected from the following groups: $-O-$; $-NA-$; $-NA_2$; $-C(O)O-$; and $-C(O)N(A)-$; wherein R^2 is defined as hereinbefore; $q = 1$ or 2 ; $m + p + q = 4$; total z per molecule is from about 3 to about 50; and X^- is an anion which is compatible with fabric softener actives and adjunct ingredients; and

6. mixtures thereof;

E. optionally, from 0 to about 15% perfume; and

F. the balance water

wherein said electrolyte and said phase stabilizer, when present, provide at least one improvement selected from: lower dilution viscosity; the same, or better, stability with less principal solvent; and/or the use of principal solvents with a ClogP outside the range of from about 0.15 to about 0.64.

26. (Amended) The composition of Claim 1 wherein said phase stabilizer is nonionic surfactant derived from saturated and/or unsaturated primary, secondary, and/or branched,